

Non-Gibrat's law in the middle scale region

Masashi Tomoyose ^{1*}, Shouji Fujimoto ^{2†} and Atushi Ishikawa ^{2‡}

¹University of the Ryukyus, 903-0213, Japan

²Kanazawa Gakuin University, 920-1392, Japan

Abstract

By using numerical simulation, we confirm that Takayasu–Sato–Takayasu (TST) model which leads Pareto's law satisfies the detailed balance under Gibrat's law. In the simulation, we take an exponential tent-shaped function as the growth rate distribution. We also numerically confirm the reflection law equivalent to the equation which gives the Pareto index μ in TST model. Moreover, we extend the model modifying the stochastic coefficient under a Non-Gibrat's law. In this model, the detailed balance is also numerically observed. The resultant pdf is power-law in the large scale Gibrat's law region, and is the log-normal distribution in the middle scale Non-Gibrat's one. These are accurately confirmed in the numerical simulation.

PACS code : 89.65.Gh

1 Introduction

Recent developments of computer technology enable vast quantities of economic data to be analyzed. As one of remarkable examples, Fujiwara et al. [1] find that Pareto's law [2]

$$P(x) \propto x^{-(\mu+1)} \quad \text{for} \quad x > x_0 \quad (1)$$

can be derived from the law of detailed balance

$$P_{12}(x_1, x_2) = P_{12}(x_2, x_1) \quad (2)$$

and Gibrat's law [3]

$$Q(R|x_1) = Q(R) , \quad (3)$$

which are observed in massive amounts of economic data. Here x is wealth, income, profits, assets, sales, the number of employees and so forth, and x_1, x_2 are two successive those. $P_{12}(x_1, x_2)$ is a joint probability density function (pdf), and $Q(R|x_1)$ is a conditional pdf of the growth rate $R = x_2/x_1$.

*e-mail address: mtomo@weiss.sci.u-ryukyu.ac.jp

†e-mail address: fujimoto@kanazawa-gu.ac.jp

‡e-mail address: ishikawa@kanazawa-gu.ac.jp

The Pareto's law (1) is not observed below some threshold x_0 [3, 4], because the Gibrat's law (3) does not hold in the middle scale region [5, 6, 7]. In Ref. [8], we show that the detailed balance (2) and a Non-Gibrat's law

$$Q(R|x_1) = \text{Const. } R^{\mp t_{\pm}(x_1)-1} \quad \text{for } R \gtrless 1 \quad (4)$$

$$t_{\pm}(x_1) = t_{\pm}(x_0) \pm \alpha \ln \frac{x_1}{x_0} \quad (5)$$

lead the log-normal distribution in the middle scale region as follows:

$$P(x_1) = C x_1^{-(\mu+1)} e^{-\alpha \ln^2 \frac{x_1}{x_0}} \quad \text{for } x_{\min} < x_1 < x_0. \quad (6)$$

The Non-Gibrat's law means a statistical dependence of the growth rate distribution on the past value x_1 in the middle scale region. In the derivation, we employ the approximation (4) observed in profits data of Japanese firms[9], and find that the expression of $t_{\pm}(x_1)$ (5) is unique under the detailed balance. Employing such empirical data, we confirm that the resultant distribution (6) fits with data in the large scale Pareto's law region and those in the middle scale log-normal region consistently. The parameters are estimated as follows: $\alpha \sim 0$ for $x_1 > x_0$, $\alpha \sim 0.14$ for $x_{\min} < x_1 < x_0$,¹ $x_0 \sim 63,000$ thousand yen and $x_{\min} \sim 1,600$ thousand yen [8].

These data analyses are sufficiently rigorous, but at the same time are restricted in the finite period and category, because it is not easy to procure exhaustive data. If we build a model based on the detailed balance and (Non-)Gibrat's law, various economic situations can be simulated. This might make the reason clear why the parameters take the empirical values. Furthermore, we can study assets or sales data of firms which are difficult to obtain. In this paper, we will propose the simulation model based on the observation of economic data.

2 TST model

Firstly, we identify the model which leads Pareto's law based on the detailed balance and Gibrat's law in the large scale region. One of the simplest and powerful candidates is Takayasu–Sato–Takayasu (TST) model [10], which is given by the Langevin equation

$$x(t+1) = b(t)x(t) + f(t), \quad (7)$$

where $b(t)$ is a non-negative stochastic coefficient and $f(t)$ is a random additive noise. They show that the conditions

$$\langle \ln b(t) \rangle < 0, \langle b(t)^2 \rangle > 1 \quad (8)$$

are necessary and sufficient for the power-law (1), the index μ of which is given by the equation

$$\langle b(t)^{\mu} \rangle = 1. \quad (9)$$

Here $\langle \dots \rangle$ denotes an average over realizations.

¹ Here, a constant parameter α takes different values in two regions. This is not an exact procedure. However, for firms which are in the large scale region in both years ($x_1 > x_0$ and $x_2 > x_0$) or in the middle scale one ($x_{\min} < x_1 < x_0$ and $x_{\min} < x_2 < x_0$), this procedure is exact. In the database, most firms stay in the same region. This parameterization is, therefore, approximately valid for describing the probability density function $P(x_1)$. This is empirically confirmed in Ref. [8].

In order to claim that this model is consistent with the empirical observation, the detailed balance and Gibrat's law must be satisfied. From Eq. (7), Gibrat's law is hold in the region where a noise $f(t)$ is negligible. The result in Ref. [1] suggests, therefore, that there is the detailed balance in TST model which leads Pareto's law. In the numerical simulation, we adopt the distribution of $b(t)$ and $f(t)$ to be Eqs. (4)–(5) with $\alpha = 0$ and Weibull distribution: $\frac{k}{\lambda}(\frac{x}{\lambda})^{k-1} \exp[-(\frac{x}{\lambda})^k]$, respectively. The typical scatter plot of the simulation is shown in Fig. 1, where we take $t_+(x_0) = 2.5$, $t_-(x_0) = 1.5$, $x_0 = 10^0$, $k = 0.5$ and $\lambda = 30$. To check the validity of the detailed balance, we take the one-dimensional Kolmogorov-Smirnov (K-S) test. We compare the distribution sample for $P(x_1 \in [10^{3+0.2(n-1)}, 10^{3+0.2n}], x_2)$ with another sample for $P(x_1, x_2 \in [10^{3+0.2(n-1)}, 10^{3+0.2n}])$ with $n = 1, 2, \dots, 20$ by making the null hypothesis that these two samples are taken from a same parent distribution. Each p value is shown in Fig. 2. The null hypothesis is not rejected in 5% significance level in the region where the noise $f(t)$ is negligible. We recognize that the detailed balance is observed in the region. The lower bound of the region is denoted by $x_{\min}$, above which Pareto's law is observed (Fig. 3).




Figure 1: The scatter plot of data points under Gibrat's law, the number of which is “500,000”.




Figure 2: Each p value of the one-dimensional Kolmogorov-Smirnov test for Fig. 1.




Figure 3: The cumulative distribution of $P(x)$ shows the Pareto's law above $x_{\min} \sim 10^4$. The Pareto index μ is estimated by nearly 1.




Figure 4: The relation between μ and $t_+(x_0) - t_-(x_0)$. Here, μ is estimated by the least square method of the simulation result, in which we take various values of $t_{\pm}(x_0)$.

In addition, we also confirm the reflection law $Q(R) = R^{-\mu-1}Q(R^{-1})$ derived in Ref. [1]. In

this simulation, from Eqs.(4) and (5) with $\alpha = 0$, the reflection law is reduced to ²

$$\mu = t_+(x_0) - t_-(x_0), \quad (10)$$

which is equivalent to the condition (9). We observe this reflection law in various $t_{\pm}(x_0)$ (Fig. 4). From these results, TST model is appropriate satisfying the detailed balance under the Gibrat's law.

3 Simulation under Non-Gibrat's law

If the detailed balance is satisfied even under a Non-Gibrat's law in the simulation, the log-normal distribution must be reduced in the middle scale region [8]. In a stochastic method, this scheme is not obvious analytically. Because a non-negative stochastic coefficient in Eq. (7) must be modified as $b(x(t), t)$ under the Non-Gibrat's law. This extension exceeds the analytical framework of TST model. However, we can examine the scheme numerically.

In the simulation, x_{int} is introduced satisfying $t_+(x_{\text{int}}) = t_-(x_{\text{int}})$. We adopt the distribution of b to be Eqs. (4)–(5) with $\alpha = 0$ for $x > x_0$ (Gibrat's law), $\alpha \neq 0$ for $x_{\text{int}} < x < x_0$ (Non-Gibrat's law) and $\alpha = 0$ for $x < x_{\text{int}}$ keeping $t_{\pm}(x)$ continuously (Fig. 5). The last parameterization is imposed to exclude immoderate slopes of growth rate distributions. As the Non-Gibrat's region in the middle scale, we observe the region $\max(x_{\text{min}}, x_{\text{int}}) < x < x_0$.




Figure 5: Continuous functions $t_{\pm}(x) = t_{\pm}(x_0) \pm \alpha \ln \frac{x}{x_0}$ with $\alpha = 0$ for $x > x_0$, $\alpha > 0$ for $x_{\text{int}} < x < x_0$ and $\alpha = 0$ for $x < x_{\text{int}}$. The horizontal axis is logarithmic scale.




Figure 6: The scatter plot of data points under Non-Gibrat's law, the number of which is “500,000 ”.

The typical scatter plot of the simulation is shown in Fig. 6, where we take $t_+(x_0) = 2.5$, $t_-(x_0) = 1.5$, $x_0 = 10^6$, $k = 0.5$, $\lambda = 30$ and $\alpha = 0.1$. To check the validity of the detailed balance, we take the same K-S test in the previous section. The result is shown in Fig. 7. The detailed balance is confirmed not only in the Gibrat's large scale region ($x > x_0 = 10^6$) but also in the Non-Gibrat's middle scale one ($\max(x_{\text{min}}, x_{\text{int}}) = 10^4 < x < x_0 = 10^6$). The resultant pdf of x is shown in Fig. 8, where the log-normal distribution in the middle scale region is observed in addition to Pareto's law in the large scale one. To confirm the validity of the log-normal distribution, we check the correlation between the parameter α inputted in the simulation and

² This equation is also confirmed empirically in profits data of Japanese firms [8].




Figure 7: The result of one-dimensional Kolmogorov-Smirnov test for Fig. 6.




Figure 8: The resultant pdf $P(x)$ with $t_+(x_0) = 2.5$, $t_-(x_0) = 1.5$, $x_0 = 10^6$, $k = 0.5$, $\lambda = 30$ and $\alpha = 0.1$. In the large scale region, Pareto's law is observed. In the middle scale one, the log-normal distribution is observed.

that estimated by fitting the pdf in the Non-Gibrat's region (denoted by α_{fit}). The result is shown in Fig. 9 for the case $t_+(x_0) = 2.5$, $t_-(x_0) = 1.5$, $x_0 = 10^6$, $k = 0.5$ and $\lambda = 30$ and in Fig. 10 for the case $t_+(x_0) = 3.5$, $t_-(x_0) = 2.0$, $x_0 = 10^6$, $k = 0.5$ and $\lambda = 30$ for example. The correlation is very high. Consequently, this simulation model is also appropriate satisfying the detailed balance even under the Non-Gibrat's law.




Figure 9: The relation between α and α_{fit} for the case $t_+(x_0) = 2.5$, $t_-(x_0) = 1.5$, $x_0 = 10^6$, $k = 0.5$ and $\lambda = 30$.




Figure 10: The relation between α and α_{fit} for the case $t_+(x_0) = 3.5$, $t_-(x_0) = 2.0$, $x_0 = 10^6$, $k = 0.5$ and $\lambda = 30$.

4 Summary and future problems

In this study, by using numerical simulation, we confirm that TST model which leads Pareto's law satisfies the detailed balance under Gibrat's law. In the simulation, we take an exponential tent-shaped function as the growth rate distribution. We also numerically confirm the reflection law [1] equivalent to the equation which gives the Pareto index μ in TST model. Moreover, we extend the model modifying the stochastic coefficient under a Non-Gibrat's law. In this model,

the detailed balance is also observed. The resultant pdf is power-law in the large scale Gibrat's law region, and is the log-normal distribution in the middle scale Non-Gibrat's one. These are accurately confirmed in the numerical simulation.

In this simulation, we employ the Non-Gibrat's law (5) that the probability of the positive growth decreases and that of the negative growth increases symmetrically as the classification of x increases (Figs. 5, 11). This is uniquely derived from the exponential tent-shaped form (4) observed in Japanese profits data [8]. On the other hand, as for sales or assets of firms, it is reported that the probability of the positive and negative growth decrease simultaneously as the classification of x increases (Fig. 12). This difference is thought to be caused by the form of the exponential tent-shaped growth rate distribution. Eq. (4) is rewritten by using the pdf $q(r|x_1)$ for $r = \log_{10} R$ as follows:

$$\log_{10} q(r|x_1) = c \mp t_{\pm}(x_1) r \quad \text{for } r \gtrless 0. \quad (11)$$

For sales or assets of firms, these linear approximations are not used to express the growth rate distribution with the curvature [1].




Figure 11: In the middle scale region, the probability of the positive growth decreases and the probability of the negative growth increases symmetrically as the classification of x increases.




Figure 12: As for sales or assets of firms, the probability of the positive and negative growth decrease simultaneously as the classification of x increases.

In order to count the curvature, we add a second order term with respect to r :

$$\log_{10} q(r|x_1) = c \mp t_{\pm}(x_1) r + \ln 10 u_{\pm}(x_1) r^2 \quad \text{for } r \gtrless 0. \quad (12)$$

In terms of $Q(R|x_1)$, these are expressed as

$$Q(R|x_1) = d R^{-1 \mp t_{\pm}(x_1) + u_{\pm}(x_1) \ln R} \quad \text{for } R \gtrless 1. \quad (13)$$

In this case, the detailed balance (2) is reduced to be

$$\frac{P(x_1)}{P(x_2)} = \frac{1}{R} \frac{Q(R^{-1}|x_2)}{Q(R|x_1)} = R^{1+t_+(x_1)-t_-(x_2)-[u_+(x_1)-u_-(x_2)] \ln R} \quad (14)$$

for $R > 1$. By setting $R = 1$ after differentiating Eq. (14) with respect to R , differential equations

are obtained. The solutions are uniquely fixed as

$$t_+(x) = t_+(x_0) + \alpha \ln \frac{x}{x_0} + \frac{\beta}{2} \ln^2 \frac{x}{x_0} + \frac{\gamma}{3} \ln^3 \frac{x}{x_0}, \quad (15)$$

$$t_-(x) = t_-(x_0) - (\alpha - \eta) \ln \frac{x}{x_0} - \frac{\beta - \delta}{2} \ln^2 \frac{x}{x_0} - \frac{\gamma}{3} \ln^3 \frac{x}{x_0}, \quad (16)$$

$$u_+(x) = u_+(x_0) - \frac{\beta + \delta}{6} \ln \frac{x}{x_0} - \frac{\gamma}{6} \ln^2 \frac{x}{x_0}, \quad (17)$$

$$u_-(x) = u_-(x_0) - \frac{\beta - 2\delta}{6} \ln \frac{x}{x_0} - \frac{\gamma}{6} \ln^2 \frac{x}{x_0}, \quad (18)$$

$$P(x) = C x^{-(\mu+1)} \exp \left[-(\alpha - \frac{\eta}{2}) \ln^2 \frac{x}{x_0} - \frac{2\beta - \delta}{6} \ln^3 \frac{x}{x_0} - \frac{\gamma}{6} \ln^4 \frac{x}{x_0} \right] \quad (19)$$

with $\eta/2 = -u_+(x_0) + u_-(x_0)$. Same solutions are obtained for $R < 1$. Here we set the parameters as follows: $\alpha, \beta, \gamma, \delta, \eta = 0$ for $x > x_0$ (Gibrat's law), $\alpha, \beta, \gamma, \delta \neq 0$ and $\eta = 0$ for $x < x_0$ (Non-Gibrat's law) because $u_{\pm}(x_0)$ are constants.

As an index of increase and decrease of the probability of the growth rate, we examine the differential coefficient at the origin: $\frac{d}{dr} \log_{10} q(r|x_1)_{r \rightarrow \pm 0} = \mp t_{\pm}(x_1)$. We are, therefore, interested in the increase and decrease of $t_{\pm}(x)$ in the middle scale region. In the case $\gamma > 0$ for example, the results are shown in Figs. 13 and 14 using the notation $X \equiv \ln(x/x_0)$, $X_{\min} \equiv \ln(x_{\min}/x_0)$. Consequently the solution, that the probability of the positive and negative growth decrease simultaneously as the classification of x increases, exists in following parameter regions:

$$\alpha = 0 \quad \text{and} \quad \delta < \beta < 0 \quad \text{and} \quad \gamma = 0, \quad (20)$$

$$\alpha = 0 \quad \text{and} \quad \delta - \gamma X_{\min} < \beta < 0 \quad \text{and} \quad \gamma > 0, \quad (21)$$

$$\alpha = 0 \quad \text{and} \quad \delta < \beta < -\gamma X_{\min} \quad \text{and} \quad \gamma < 0. \quad (22)$$

In any case, $t_{\pm}(x)$ have no first order term and the negative second order term with respect to $\ln(x/x_0)$. Not only parameter β but also δ is necessary for the solution. We will apply the simulation to this analysis in the next work.




Figure 13: For $\gamma > 0$, the monotonically increasing or decreasing regions of $t_+(x)$.




Figure 14: For $\gamma > 0$, the monotonically increasing or decreasing regions of $t_-(x)$.

As other application of the simulation, the detailed balance is expected to be extended to the detailed quasi-balance [11] observed in the assessed value of land in Japan [12, 13]. If these applications are studied adequately, we can manage the risk of these quantities by using knowledge of this method.

Acknowledgements

We would like to thank the Yukawa Institute for Theoretical Physics at Kyoto University, where this work was initiated during the YITP-W-07-16 on “Econophysics III–Physical Approach to Social and Economic Phenomena–”, and especially to Dr Y. Soejima for the important comments about our work. Thanks are also due to Dr. A. Sato for a lot of useful discussions. This work was supported in part by a Grant-in-Aid for Scientific Research (C) (No. 20510147) from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

References

- [1] Y. Fujiwara, W. Souma, H. Aoyama, T. Kaizoji and M. Aoki, *Physica* **A321** (2003), 59.
Y. Fujiwara, C.D. Guilmi, H. Aoyama, M. Gallegati and W. Souma, *Physica* **A335** (2004), 197.
- [2] V. Pareto, *Cours d'Economie Politique* (Macmillan, London, 1897).
- [3] R. Gibrat, *Les inegalites economiques* (Sirey, Paris, 1932).
- [4] W.W. Badger, in: *B.J. West (Ed.), Mathematical Models as a Tool for the Social Science* (Gordon and Breach, New York, 1980), p. 87.
E.W. Montrill, M.F. Shlesinger, *J. Stat. Phys.* **32** (1983), 209.
- [5] M.H.R. Stanley, L.A.N. Amaral, S.V. Buldyrev, S. Havlin, H. Leschhorn, P. Maass, M.A. Salinger, H.E. Stanley, *Nature* **379** (1996), 804.
- [6] H. Takayasu, M. Takayasu, M.P. Okazaki, K. Marumo, T. Shimizu, cond-mat/0008057, in: *M.M. Novak (Ed.), Paradigms of Complexity* (World Scientific, 2000), p. 243.
- [7] H. Aoyama, Ninth Annual Workshop on Economic Heterogeneous Interacting Agents (WE-HIA 2004).
H. Aoyama, Y. Fujiwara, W. Souma, The Physical Society of Japan 2004 Autumn Meeting.
- [8] A. Ishikawa, *Physica* **A367** (2006), 425; *Physica* **A383/1** (2007), 79.
- [9] TOKYO SHOKO RESEARCH, LTD., <http://www.tsr-net.co.jp/>.
- [10] H. Takayasu, A. Sato and M. Takayasu, *Phys. Rev. Lett.* **79** (1997), 966.
- [11] A. Ishikawa, *Physica* **A371** (2006), 525.
- [12] T. Kaizoji, *Physica* **A326** (2003), 256.
T. Kaizoji and M. Kaizoji, *Physica* **A344** (2004), 138.
- [13] The Ministry of Land, Infrastructure and Transport Government of Japan's World-Wide Web site, <http://nlftp.mlit.go.jp/ksj/>.